

Reduction of CuO in H₂: In Situ Time-Resolved XRD Studies

J. Rodriguez, J. Kim, J. Hanson (BNL), M. Pérez (Univ. Central Venezuela), A. Frenkel (Yeshiva)
Beamline(s): X7B

Introduction: CuO is used as a catalyst or catalyst precursor in many chemical reactions that involve hydrogen as a reactant or product. It has been proposed that in several of these catalytic process CuO undergoes a complete reduction and metallic copper(Cu⁰) is the real active phase [1]. For years, there has been controversy about the relative importance of Cu⁺¹ and Cu⁰ in centers in the methanol synthesis reaction [2,3]. In order to solve this issue one needs a fundamental understanding of the H₂ reaction with CuO [4].

Methods and Materials: Samples of CuO were loaded in a sapphire capillary attached to a flow-reaction cell similar to those described in refs. [5,6]. The sample was heated in the range of 150-300°C with a small resistance heater wrapped around the capillary. A 5%H₂/95%He mixture was flowed through the capillary and reactant gases were sampled with a SRS RGA. The XRD data were collected with a MAR345 image plate detector and the powder rings were integrated using FIT2D code [7].

Results: Fig. 1 shows typical results from the time resolved XRD and Fig. 2 shows H₂O formation determined from RGA.

Conclusions: Oxide reduction was observed but only after an induction period. High temperature or increased H₂ pressure lead to a decrease in the magnitude of the induction time. The H₂ flow rate determines the amount of Cu⁺¹ formation. Under higher flow rates a direct CuO → Cu transformation occurs. To facilitate the generation of Cu⁺¹ in a catalytic process one can limit the supply of H₂ or mix this molecule with molecules which can act as oxidizing agents (O₂, H₂O).

Acknowledgments: The work at BNL was financed through contract De-AC02-98CH10086 with the US DOE Office of Basic Energy Sciences, Division of Chemical Sciences.

References: [1]. Campbell, C. T., Daube, K. A. and White, J.M. *Surf. Sci.*, **182**, 458 (1987).

[2] Kung, H.H., *Transition Metal Oxides: Surface Chemistry and Catalysis* Elsevier: NewYork, 1989.

[3] Klier, K., *Adv. Catal.* **31**, 243 (1983).

[4]. Rodriguez, J. A., Kim, J.Y., Hanson, J.C., Perez, M., Frenkel, A. F. *Cat. Letters* (in press)

[5]. Chupas, P. J., Ciruolo, M.F., Hanson, J.C. and Grey, C.P., *J. Am. Chem.Soc.*, **123**, 1694 (2001).

[6]. Clausen, B. S., Steffensen, G., Fabius, B., Villadsen, J., Freidenhans, R and Topsoe, H, *J. Catal.* **132**, 524 (1991).

[7] Hammersley, A. P., Svensson, S.O. and Thompson, A., *Nucl. Instr. Methods Phys.*, **346**, 321 (1994)




Fig1. In situ time-resolved XRD reduction of CuO to Cu⁰.
(flow rate 20cc/min)




Fig. 2 Water formation during reduction in Fig. 1.